Alpine Entomology 8 2024, 35–49 | DOI 10.3897/alpento.8.117537

PENSOFT

Alpine Entomology

Distribution of the invasive ambrosia beetle *Anisandrus maiche* (Coleoptera, Scolytinae) in Switzerland and first record in Europe of its ambrosia fungus *Ambrosiella cleistominuta*

José P. Ribeiro-Correia[1,2], Simone Prospero[1], Ludwig Beenken[1], Peter H. W. Biedermann[3], Simon Blaser[1], Manuela Branco[2], Yannick Chittaro[4], David Frey[5], Doris Hölling[1], Sezer Olivia Kaya[1,6], Miloš Knížek[7], Jana Mittelstrass[1], Beat Ruffner[1], Andreas Sanchez[4], Eckehard G. Brockerhoff[1]

1 *Swiss Federal Institute for Forest, Snow and Landscape Research WSL, CH-8903 Birmensdorf, Switzerland*

2 *Universidade de Lisboa, Instituto Superior de Agronomia, Centro de Estudos Florestais Tapada da Ajuda, PT-1349-017 Lisboa, Portugal*

3 *Chair of Forest Entomology and Protection, Institute of Forestry, University of Freiburg, Fohrenbühl 27, D-79252 Stegen-Wittental, Germany*

4 *info fauna, Avenue Bellevaux 51, CH-2000 Neuchâtel, Switzerland*

5 *Al Ciòs Consulenze ambientali, Via Cantonale 79, CH-6818 Melano, Switzerland*

6 *Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences, Box 190, 23422 Lomma, Sweden*

7 *Forestry and Game Management Research Institute, Strnady 136, CZ-252 02 Jíloviště, Czech Republic*

https://zoobank.org/6AC85CAE-03C7-4137-8069-97A437ECD326

Corresponding authors: Eckehard G. Brockerhoff (eckehard.brockerhoff@wsl.ch); José P. Ribeiro-Correia (ribeiro.correia.jp@gmail.com)

Academic editor: Christoph Germann ♦ **Received** 18 December 2023 ♦ **Accepted** 22 February 2024 ♦ **Published** 18 March 2024

Abstract

Ambrosia beetles are highly successful as invaders because they are often transported internationally with wood packaging and other wood products and because their inbreeding mating systems facilitates establishment of invading populations. In 2022, two independent insect surveys in canton Ticino (southern Switzerland) revealed the widespread occurrence of the invasive ambrosia beetle *Anisandrus maiche* (Kurentzov, 1941) from southern to central-upper Ticino. This species is native to east Asia and has previously been found as a non-native invasive species in the United States, Canada, western Russia, Ukraine and, in 2021, in northern Italy. Here, we present the results of several trapping studies using different trap types (bottle traps, funnel traps and Polytrap intercept traps) and attractants and a map of the distribution of the species. In total, 715 specimens of *A. maiche*, all female, were trapped, and the identity of selected individuals was confirmed by morphological and molecular identification based on three mitochondrial and nuclear markers (COI, 28S and CAD). Trap samples from early April to early September 2022 in intervals of two to four weeks showed that flights of *A. maiche* occurred mainly from June to mid-August. Isolation of fungal associates of *A. maiche* from beetles trapped alive revealed the presence of four fungal species, including the ambrosia fungus *Ambrosiella cleistominuta*, the known mutualist of *A. maiche*. The identity of *A. cleistominuta* was confirmed by comparing DNA sequences of its nuclear, internal transcribed spacer (ITS) gene with reference sequences in NCBI and BOLDSYSTEMS. This represents the first record of *A. cleistominuta* in Europe. Of the other fungal associates isolated from *A. maiche* in Ticino, *Fusarium lateritium* is of note as there is a possibility that *A. maiche* could act as a vector of this plant pathogen. We highlight several research needs that should be addressed to gain insight into the potential impact of these non-native species and to overcome problems with heteroplasmy in COI sequences in studies of invasion and population genetics of ambrosia beetles.

Key Words

Bark and ambrosia beetles, biological invasions, Ceratocystidaceae, Curculionidae, detection, surveillance

Copyright *José P. Ribeiro-Correia et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Biological invasions are a growing concern due to the continuing increase in establishments of non-native (alien) invasive species and their impacts on native species, natural and modified ecosystems and on plant, animal and human health (Kenis et al. 2009; Brockerhoff et al. 2017; Seebens et al. 2018; Pyšek et al. 2020). Bark and ambrosia beetles (Coleoptera: Curculionidae, Scolytinae) are particularly successful as invaders because they are often transported internationally by trade and transport of wood products and due to the widespread use of wood packaging materials such as pallets and crates (Brockerhoff et al. 2006; Lantschner et al. 2020). In addition, many ambrosia beetles have mating systems that involve inbreeding (Kirkendall 1983) which greatly facilitates the establishment of new populations by very few invading individuals (Kirkendall and Faccoli 2010; Lantschner et al. 2020). Furthermore, because ambrosia beetles feed only on specific ambrosia fungi (on which they are nutritionally dependent and which they carry in their mycangia and cultivate in galleries they excavate in deadwood), these beetles often have a very wide host range (e.g., Ranger et al. 2016; Hulcr and Stelinski 2017). This also contributes to their invasiveness since they are less dependent on the presence of a particular host species. Native ambrosia fungi are typically not pathogenic, but some non-native ambrosia beetles are vectors of severe plant pathogens that can cause tree death. For example, the Asian *Xyleborus glabratus* Eichhoff, 1877 and its symbiotic tree pathogen *Raffaelea lauricola* T.C. Harr., Fraedrich & Aghayeva (Harrington et al. 2008), the causal agents of laurel wilt, are responsible for large-scale mortality of many trees in the Lauraceae family in the southeastern United States (Fraedrich et al. 2008; Hughes et al. 2017).

Anisandrus maiche (Kurentzov, 1941) is an invasive ambrosia beetle native to northeast Asia (i.e., parts of China, Japan, North and South Korea, and the Russian Far East) (Knížek 2011; Terekhova and Skrylnik 2012; Alonso-Zarazaga et al. 2017, 2023; Park et al. 2020; Smith et al. 2020; EPPO 2022). Established non-native populations of *A. maiche* were detected in North America first in 2005 in Pennsylvania, USA, and subsequently in the US states of Ohio, West Virginia, Illinois, Indiana, Maryland, New Jersey, New York, and Wisconsin, and in the Canadian Provinces Ontario and Quebec (Rabaglia et al. 2009; Haack et al. 2013; Gomez et al. 2018; Thurston et al. 2022). In Europe, non-native populations were detected first in 2007 in western Russia and in eastern Ukraine (Moscow oblast, Belgorod oblast, Donetsk oblast, Kharkiv oblast and Sumy oblast (Nikulina et al. 2007; Nikitskii 2009; Terekhova and Skrylnik 2012; Kovalenko and Nikitski 2013; Nikulina et al. 2015)). About 14 years later, in 2021–2022, this beetle was found in the Italian regions Veneto and Lombardy (Colombari et al. 2022; Ruzzier et al. 2022).

In July 2022, we collected several specimens of *A. maiche* in trap catches from two locations in central canton Ticino, in southern Switzerland. Subsequently, many more specimens were found in trap catches from several other locations across the southern half of canton Ticino and in the southwestern part of canton Grisons, suggesting that the species has been established in these areas for several years. As of 2023, *Anisandrus maiche* is regulated as a quarantine pest in Switzerland (PGesV (2018) and PGesV-WBF-UVEK (2019)) and the EU, as are all non-European Scolytinae that are not present in the EU. However, its classification as a quarantine pest is being reviewed.

Here, our objectives are to report the discovery of *Anisandrus maiche* in Switzerland, the locations and forest types where it was found, and the traps and attractants with which the species was caught. Furthermore, we provide information on the fungal and microbial associates which we recorded from *A. maiche* in Switzerland, including its ambrosia fungus *Ambrosiella cleistominuta* C. Mayers & T.C. Harr., and on potential damage caused by this beetle, based on a review of available information.

Materials and methods

Abbreviations used

FPS Forest Protection Switzerland (Waldschutz Schweiz), Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland;

MCSN Museo cantonale di storia naturale (Natural History Museum), Lugano, Switzerland;

PHP Phytopathology, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland;

UPN Ufficio della Natura e del Paesaggio, Bellinzona, Switzerland;

WSL Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland

Study sites of *A. maiche* in cantons Ticino and Grisons, and description of traps and attractants

All study sites in Switzerland where *A. maiche* was found are located in the southern part of the country in the cantons Ticino (40 sites) and Grisons (one site, Roveredo) (Table 1, Fig. 1). The sites were in mixed forests composed mostly of sweet chestnut (*Castanea sativa*), oak (*Quercus* sp.), Scots pine (*Pinus sylvestris*) and other tree species, with varying proportions of broadleaved trees and conifers.

Five trap type and lure combinations were used in 2022 during several insect surveys in which *A. maiche*

Table 1. Study site locations, traps used and the number of *Anisandrus maiche* trapped in 2022. Trap types: BT, bottle trap; FG, funnel trap green; FB, funnel trap black; PT, Polytrap. Lures: E, ethanol; EA, ethanol + alpha-pinene; EA8, ethanol + alpha-pinene + eight-component blend (for details about trap types and lures see methods).

Study site location and Canton*	Swiss coordinates (E, N (LV 95))	Degrees latitude, longitude (WGS84)	Elevation (m a.s.l.)	Trap type and lure	Number of *A. maiche* trapped (total / number per trap day from June to August)
(1) Riviera-Iragna 1, TI	2717293, 1133879	46.3460, 8.9623	286	BT/E	134 / 1.47
(2) Riviera-Iragna 2, TI	2717138, 1134143	46.3484, 8.9604	285	BT/E	50 / 0.53
(3) Serravalle-Leggiuna 1, TI	2718898, 1138748	46.3895, 8.9844	362	BT/E	53 / 0.51
(4) Serravalle-Leggiuna 2, TI	2718937, 1138900	46.3909, 8.9849	386	BT/E	11 / 0.16
(5) Faido-Anzonico 1, TI	2708760, 1144009	46.4386, 8.8539	970	BT/E	0 / 0.00
(6) Faido-Anzonico 2, TI	2708628, 1143968	46.4382, 8.8521	916	BT/E	0 / 0.00
(7) Brusino-Arsizio, TI	2715995, 1086645	45.9215, 8.9338	320	BT/E	36 / 1.20
(8) Bellinzona-Camorino, TI	2722160, 1113855	46.1651, 9.0203	460	BT/E	23 / 0.77
(9) Losone, TI	2700175, 1114515	46.1747, 8.7359	275	BT/E	111 / 3.70
(10) Novazzano, TI	2720790, 1078230	45.8449, 8.9934	268	BT/E	45 / 1.50
(11) Roveredo, GR	2730490, 1120800	46.2260, 9.1301	505	BT/E	41 / 1.37
(12) Rivera-Carona, TI	2714268, 1110343	46.1349, 8.9173	517	FG/EA8	5 / 0.06
(13) Moleno, TI	2720086, 1125929	46.2741, 8.9966	255	FG/EA8	6 / 0.07
(14) Chiasso-Bresciano, TI	2721348, 1077091	45.8346, 9.0003	262	FG/EA8	1 / 0.01
(15) Riviera-Iragna 3, TI	2717083, 1134188	46.3489, 8.9597	286	FB/EA8	1 / 0.01
(16) Chiasso-Pian Pessina, TI	2723687, 1076070	45.8250, 9.0302	440	FB/EA8	0 / 0.00
(17) Chiasso-Bresciano, TI	2721348, 1077091	45.8346, 9.0003	262	FB/EA8	2 / 0.02
(18) Faido-Anzonico 1, TI	2708760, 1144020	46.4387, 8.8539	970	FB/EA	0 / 0.00
(19) Faido-Anzonico 2, TI	2708628, 1143968	46.4382, 8.8522	915	FB/EA	2 / 0.02
(20–134) entire Canton TI	See Sanchez et al. (2023) and Suppl. material 1			PT / –	194 / 0.07

* Two-letter abbreviation for Cantons, TI – Ticino, GR – Grisons.



Figure 1. Trap locations in cantons Ticino and Grisons where *Anisandrus maiche* was captured (green symbols) or where no captures were recorded (grey symbols). Symbols vary by trap type (see legend and methods for details). Note that each square represents a pair of two Polytraps which were placed in close proximity to each other (Vector and raster map data https://www.swisstopo.ch).

was caught. **Type 1** (**bottle traps**, "**BT**") traps consisted of bottle traps made at the laboratory based on the design described in Grégoire et al. (2001) and as modified by Gossner et al. (2019) with 70% ethanol as lure and propylene glycol as preservative and installed such that the bottle part was about 1 m above ground (Suppl. material 3: fig. 1a). Two BT were installed at three study sites near Biasca (see Table 1) on 2 April 2022 and taken down on 26 August 2022. Five further single BT were installed at additional study sites (see Table 1) on 18 July 2022 and removed on 30 August 2022. A small number of additional "live bottle traps" were used without propylene glycol to enable live captures for studies of fungal associates (see below).

Type 2 (**funnel traps green**, "**FG**") traps were green multi-funnel traps (ChemTica Internacional, Costa-Rica, Suppl. material 3: fig. 1b) with propylene glycol as preservative and baited with alpha-pinene and ethanol (Econex, Spain) and an 8-component lure blend (fuscumol, fuscumol acetate, geranylacetone, monochamol, 3-hydroxyhexan-2-one, anti-2,3-hexanediol, 2-methylbutan-1-ol, and prionic acid) as described by Fan et al. (2019). FG traps were suspended by a rope inside the canopy of trees at a height of approximately 10 m. One of these traps was installed at each study site (see Table 1) on 12 April 2021 and taken down on 1 or 29 September 2021. These traps, as well as the black funnel traps mentioned below, were part of a separate project aimed at early detection of quarantine pests, especially longhorned beetles (Cerambycidae) but they are also effective for bark and ambrosia beetles (see Roques et al. 2023).

Type 3 (**funnel traps, black**, "**FB**") traps were black multi-funnel traps (ChemTica Internacional, Costa-Rica, Suppl. material 3: fig. 1c) with propylene glycol as preservative and the same lure combination and trapping period as Type 2. FB traps were suspended by a rope below the canopy of trees at a height of 2–3 m above ground. One of these traps was installed at each study site (see Table 1) on 12 April 2021 and taken down on 1 or 29 September 2021. Trapping with funnel traps (FG and FB) in 2021 was undertaken as part of a surveillance programme aimed at priority quarantine pests targeting mainly longhorn beetles.

Type 4 (**Polytrap**, "**PT**") traps were unbaited Polytrap interception traps (as described by Brustel 2012, Suppl. material 3: fig. 1d) with saturated salt solution and neutral detergent as preservative, suspended 2 m above the ground. As part of a biodiversity survey funded by canton Ticino (MCSN and UPN), a total of 114 Polytraps were installed on 10 March 2022 and taken down at the end of 3 October 2022 (see Sanchez et al. (2023) for details).

In addition to the traps described above, collection of ambrosia beetles was also attempted with log sections of European beech (*Fagus sylvatica*), Norway spruce (*Picea abies*) and sweet chestnut ca. 50 cm long, 5–10 cm diameter, baited with 70% ethanol and suspended alongside tree stems at a height of 1.5–2.0 m, as described by Monterrosa et al. (2021). These were installed at study sites (1)–(6) in Ticino (Table 1) from 16 April to 26 August 2022. In addition, short sections of beech branches (about 20 cm long, 2–4 cm diameter) soaked in 70% ethanol were placed on the ground at study sites (1)–(6) in Ticino from 12 June to 25 July 2022 for two periods of approximately three weeks.

Study sites in cantons Valais and Zurich

In addition to the traps placed in cantons Ticino and Grisons, bottle traps with ethanol as lure (as trap type 1 described above) were also used in canton Valais from 18 March 2022 to 25 August 2022 at six locations (Brig 1, 46.2905°N, 7.9601°E, 1260 m a.s.l.; Brig 2, 46.2941°N, 7.9577°E, 1259 m a.s.l.; Lens 1, 46.2677°N, 7.4339°E, 1096 m a.s.l.; Lens 2, 46.2677°N, 7.4339°E, 1097 m a.s.l.; Visp 1, 46.2971°N, 7.8566°E, 676 m a.s.l.; and Visp 2, 46.2967°N, 7.8564°E, 705 m a.s.l.), and in canton Zurich from 4 April 2022 to 31 August 2022 at six locations (Zurich-Hönggerberg 1, 47.4121°N, 8.4978°E, 535 m a.s.l.; Zurich Hönggerberg 2, 47.4196°N, 8.4872°E, 525 m a.s.l.; Stallikon-Uetliberg 1, 47.3364°N, 8.4936°E, 660 m a.s.l.; Stallikon-Uetliberg 2, 47.3367°N, 8.4945°E, 669 m a.s.l.; Birmensdorf-Rameren 1, 47.363°N, 8.4483°E, 540 m a.s.l.; Birmensdorf Rameren 2, 47.3631°N, 8.4483°E, 555 m a.s.l.). The study sites in cantons Valais and Zurich were also in mixed forests with varying proportions of broadleaved trees and conifers and composed mostly of oaks, beech, Norway spruce, Scots pine or other trees.

Curation and morphological and molecular identification of *A. maiche*

All ambrosia and bark beetles from traps with preservative were sorted under a stereomicroscope and kept in 70% ethanol for temporary storage while selected individuals were point mounted. Ambrosia and bark beetles were identified morphologically by J. Ribeiro-Correia, E. Brockerhoff, A. Sanchez and M. Knížek using Grüne (1979), Pfeffer (1995), Stark (1952), Rabaglia et al. (2009) and reference collections held at WSL, by M. Knížek and by Andreas Sanchez. A specimen of *A. maiche* from Serravalle-Leggiuna 1 is shown in Fig. 2.

For molecular diagnostics, genomic DNA from *A. maiche* was extracted from adults using the NucleoSpin Tissue XS Kit (Macherey-Nagel, Düren, Germany) using whole insects or, to preserve the specimens, leg fragments and one elytron. The COI barcode region was amplified and sequenced with primers LCO1490/HCO2198 (Folmer et al. 1994). The nuclear markers, the ribosomal-encoding gene 28S and the CAD gene were targeted with primers 3690s/a4285 (Kambestad et al. 2017) and CADforB2/apCADrevlmod (Dole et al. 2010), respectively. PCR conditions were set as specified in the respective reference papers. Sanger sequencing was performed using the same primers as for PCR and the BigDye Terminator

Figure 2. *Anisandrus maiche* adult trapped at Serravalle-Leggiuna 1 (Ticino), dorsal and lateral. Specimen length 2.0 mm. Photos by Carl-Michael Anderson, WSL.

v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, California, USA). Sequences were trimmed and assembled using the CLC Main Workbench Version 22.0.2 (QIAGEN, Aarhus, Denmark) and checked manually before being subjected to BLAST searches against the All Barcode database on BOLD (https://www.boldsystems.org) and NCBI (https://blast.ncbi.nlm.nih.gov). All COI barcode sequences from *A. maiche* generated in this study are deposited on BOLD (accession numbers listed below). In addition, a specimen of *A. maiche* (BOLD Sequence ID: SCOL295-12) collected in Primorsky Krai, Russian Far East, was obtained from Bjarte Jordal (University of Bergen, Natural History Collections) and processed for molecular analyses as described above for comparison with our specimens.

Voucher specimens are held at WSL, Birmensdorf, Switzerland, at the MCSN, Lugano, Switzerland, and in the collections of M. Knížek in Prague, Czechia, A. Sanchez in Sion, Switzerland, and Heiko Gebhardt in Tübingen, Germany. Details on individual reference specimens are provided below.

Isolation, cultivation, and identification of fungal species associated with *A. maiche*

Live bottle traps (as described above for trap type 1, but with moistened sterile paper instead of propylene glycol in the collection jar) were installed at six locations in Ticino (locations number (1)–(6), see Table 1) to collect live beetles for isolation and cultivation of associated fungi. Traps were inspected after 2 to 4 days in the field. Beetles present in the traps were removed immediately from the PVC bottle, placed individually in 1.5 mL Eppendorf tubes and, upon arrival at the laboratory, stored at 4 °C for a maximum of one week.

The fungal species present on the surface and inside the collected beetles were identified as follows. First, the individual beetles were taken from the Eppendorf tubes using sterile tweezers and gently placed onto weaker-strength agar medium (SMA; 10g/L malt extract; 15g/L agar; 100 ppm streptomycin added after autoclaving to prevent growth of bacteria). The beetles were allowed to walk freely for 30–45 min so that fungal spores present on their body would eventually deposit on the agar surface. To identify fungal species inside their body (i.e., in the mycangia and in the digestive tract), beetles were subsequently removed from the Petri dish using sterile tweezers, placed in 90% ethanol for 1–2 seconds to kill any spores still present externally on their body, rinsed twice in sterile distilled water, and placed on a sterile paper towel to dry. Once dry, beetles were placed individually in a new 1.5 mL Eppendorf tube containing 0.5 mL distilled sterile water and crushed with a sterile rod. After brief vortexing, 100 µL of this solution was spread on SMA and incubated in the dark at room temperature. Plates were checked daily for up to one week and growing fungal colonies were subcultured on Potato Dextrose Agar (PDA; 39 g l^{-1}, Difco, Voight Global Distribution, Lawrence, MD, USA). When morphologically different colonies were present on a plate, a representative colony of each morphotype was transferred to PDA. After incubation of the PDA plates for two weeks in the dark at room temperature, fungal cultures were grouped into morphotypes based on the macro-morphological features of their mycelia.

For species identification, DNA was extracted from 1–3 representative cultures of each morphotype using LGC reagents and Kingfisher 96/Flex (LGC Genomics GmbH, Berlin, Germany), according to the manufacturer's instructions. The nuclear, internal transcribed spacer (ITS) was then amplified by PCR and sequenced in both directions using the forward ITS5 and reverse ITS4 primers (White et al. 1990) following the general methodology described in Franić et al. (2019). Sequences were assembled and edited using CLC Main Workbench Version 22.0.2 and compared with reference sequences in NCBI and BOLDSYSTEMS databases. Sequences were considered to belong to the same species if they showed at least 99% similarity.

Results

Specimens of *A. maiche* trapped in Switzerland

In 2022, a total of 715 specimens of *A. maiche* were trapped in southern Switzerland (Table 1, Fig. 1), including 504 in bottle trap samples and 194 in Polytrap samples. The first specimens were discovered in bottle trap catches from May 2022 from the vicinity of Biasca (Riviera-Iragna and Serravalle-Leggiuna), which were identified by Miloš Knížek on 1 July 2022 and confirmed by molecular analysis soon after (see below). Over the summer of 2022, 248 specimens of *A. maiche* were trapped with bottle traps (all with ethanol as lure) at Riviera-Iragna and Serravalle-Leggiuna (Table 1). An additional 256 *A. maiche* were trapped at five locations in southern Ticino with additional traps placed there to determine the extent of the distribution of *A. maiche* in Ticino (Table 1).

Two more specimens of *A. maiche* were found in samples from a black funnel trap from Faido-Anzonico in northern Ticino, the northernmost occurrence of the species (Table 1). These traps were part of another project which used funnel traps similar to trap type 3 (FB) but baited only with ethanol and alpha-pinene lures and suspended from pine branches about 2–3 m above ground.

Polytraps (trap type 4) installed in 2022 across the canton of Ticino caught 194 specimens of *A. maiche*. These were captured with 25 traps distributed between Chiasso (southernmost Ticino) and Biasca (central-upper Ticino) out of a total of 114 polytraps placed across Ticino. No specimens were captured in the Polytraps located north of Biasca and in the Maggia Valley north of Terre di Pedemonte (near Ascona) (Fig. 1).

Following the discovery of *A. maiche* in the vicinity of Biasca in June–July 2022, samples collected in a Swiss surveillance programme in 2021 (using green and black funnel traps, trap types 2 and 3) aimed at detecting priority quarantine insects, especially longhorn beetles (Cerambycidae), were re-examined for the presence of *A. maiche*. In these samples, a total of 15 *A. maiche* were found in southern Ticino near the Italian border (two sites near Chiasso), north of Lugano (Rivera), north of Bellinzona (Moleno), and near Biasca (Riviera-Iragna) (Table 1, Fig. 1).

Across all sites and trap types, most specimens of *A. maiche* were trapped at lower elevations in the valleys or lower mountain slopes at elevations between 195 m a.s.l. (near Locarno) and 386 m a.s.l. (near Biasca), but a few were caught at higher elevations such 626 m a.s.l. (Capriasca, north of Lugano) and 916 m a.s.l. (Faido Anzonico). All specimens were trapped in a variety of forest types with sweet chestnut, beech, mixed broadleaved trees and, in a few cases, a mixture with Scots pine.

No *A. maiche* were caught with bottle traps at six sites in canton Valais and six sites in canton Zurich (that were part of the same study using bottle traps described here).

Molecular identification of *A. maiche*

To confirm the identity of selected specimens of *A. maiche*, nucleotide BLAST searches were performed on BOLD and NCBI (as accessed in October 2022). For the mitochondrial COI barcode region, seven out of our 14 samples share 100% identity with accession MN619845 on NCBI, designated as *A. maiche*. For three other samples from Switzerland, a second haplotype was identified sharing 100% identity with a private accession on BOLD, also designated as *A. maiche*. The two haplotypes show a divergence of 5.9% resulting from a remarkable number of base substitutions between the haplotypes. In addition, four specimens show a pattern of heteroplasmy, compatible with the two haplotypes identified. However, assessing the nuclear, ribosomal-encoding gene 28S revealed 100% identity for all specimens to the ribosomal-encoding gene 28S of *A. maiche* (GenBank Accession MK098863, voucher specimen UFIFAS UFFE 28176). In addition, the CAD fragment from three samples (PHP22_0410, PH22_0411, PHP22_0539) displayed 100% identity to a sequence of *A. maiche* (GenBank Accession MN260139, Shanghai, China), but differed by 1bp to accession MN260138 collected in Michigan, USA . The *A. maiche* specimen from the Russian Far East (BOLD Sequence ID: SCOL295-12), which was re-analysed together with our specimens from Switzerland, showed the same pattern of COI heteroplasmy and shared 100% identity to the 28S locus of all our analysed specimens.

List of selected specimens held in collections

Selected specimens from Switzerland, collected by José Ribeiro-Correia, caught in bottle traps with ethanol as lure (see Suppl. material 1 for a complete list):

- 2 females, Leggiuna 1 (Serravalle-Leggiuna), Ticino, LV95: 2718898 E, 1138748 N (46.3895°N, 8.9844°E), 370 m a.s.l., 13–27 May 2022, det. Miloš Knížek, WSL DNA-IDs PHP22_0410 (GenBank: OQ685554) and PHP22_0411 (GenBank: OQ685554), entire specimens used for molecular analysis, DNA held at WSL PHP.
- 1 female, Iragna 1 (Riviera-Iragna), Ticino, LV95: 2717293 E, 1133879 N (46.3460°N, 8.9623°E), 286 m a.s.l., 27 May–10 June 2022, det. Miloš Knížek, WSL DNA-ID PHP22_0539 (GenBank: OQ685552), entire specimens used for molecular analysis, DNA held at WSL PHP.
- 7 females, Iragna 2 (Riviera-Iragna), Ticino, LV95: 2717138 E, 1134143 N (46.3484°N, 8.9604°E), 285 m a.s.l., 10–25 June 2022, det. José Ribeiro-Correia, WSL DNA-IDs PHP22_0627 (GenBank: OQ685551), PHP22_0628 (GenBank: OQ685550), PHP22_0629 (GenBank: OQ685549), PHP22_0630 (GenBank: OQ685548), PHP22_0946 (GenBank: OQ685547), PHP22_0947(GenBank: OQ685546),

PHP22_0952(GenBank: OQ685541), specimens at WSL FPS (ethanol/freezer), DNA held at WSL PHP.

- 1 female, Brusino-Arsizio, Ticino, LV95: 2715995 E, 1086645 N (45.9215°N, 8.9338°E), 320 m a.s.l., 18 July–3 Aug. 2022, det. José Ribeiro-Correia, WSL DNA-ID PHP22_0948 (GenBank: OQ685545), specimen at WSL FPS (ethanol/freezer), DNA held at WSL PHP.
- 3 females, Roveredo, Grisons, LV95: 2730490 E, 1120800 N, (46.2260°N, 9.1301°E), 505 m a.s.l., 18 July–3 Aug. 2022, det. José Ribeiro-Correia, WSL DNA-IDs PHP22_0949 (GenBank: OQ685544), PHP22_0950 (GenBank: OQ685543), PHP22_0951 (GenBank: OQ685542), specimens at WSL FPS (ethanol/freezer), DNA held at WSL PHP.
- 2 females, Leggiuna 1 (Serravalle-Leggiuna), Ticino, LV95: 2718898 E, 1138748 N (46.3895°N, 8.9844°E), 370 m a.s.l., 20 April–13 May 2022 / 13–27 May 2022, det. Miloš Knížek, specimens at WSL FPS (pinned).
- 3 females, Leggiuna 1 (Serravalle-Leggiuna), Ticino, LV95: 2718898 E, 1138748 N (46.3895°N, 8.9844°E), 370 m a.s.l., 13–27 May 2022, det. José Ribeiro-Correia, specimens at WSL FPS (pinned).
- 2 females, Iragna 2 (Riviera-Iragna), Ticino, LV95: 2717138 E, 1134143 N (46.3484°N, 8.9604°E), 285 m a.s.l., 13–27 May 2022, det. Miloš Knížek, specimens at Miloš Knížek collection, Prague, Czechia.
- 11 females, Iragna 1 (Riviera-Iragna), Ticino, LV95: 2717293 E, 1133879 N (46.3460°N, 8.9623°E), 286 m a.s.l., 27 May–10 June 2022, det. Miloš Knížek, specimens at Miloš Knížek collection, Prague, Czechia.
- 2 females, Leggiuna 1 (Serravalle-Leggiuna), Ticino, LV95: 2718898 E, 1138748 N (46.3895°N, 8.9844°E), 370 m a.s.l., 13–27 May 2022, det. Miloš Knížek, specimens at Miloš Knížek collection, Prague, Czechia.

Selected specimens from Switzerland, collected by David Frey, caught with unbaited Polytrap interception traps (see Suppl. material 2 for a complete list):

- 13 females, Ruderi del Castello di Claro (Bellinzona), Ticino, LV95: 2722751 E, 1124011 N (46.2563°N, 9.0306°E), 437 m a.s.l., 14–29 June 2022, det. Andreas Sanchez, specimens at MCSN.
- 11 females, Ciossa Antognini (Cadenazzo), Ticino, LV95: 2714516 E, 1113409 N (46.1624°N, 8.9212°E), 201 m a.s.l., 16 June–15 July 2022, det. Andreas Sanchez, specimens at MCSN.
- 1 female, El Piágn (Cugnasco-Gerra), Ticino, LV95: 2712487 E, 1116016 N (46,1862°N, 8,8956°E), 774 m a.s.l., 31 May–16 June 2022, det. Andreas Sanchez, specimen at MCSN.
- 1 female, Solorónch (Capriasca), Ticino, LV95: 2717651 E, 1101409 N (46.0539°N, 8.9588°E), 626 m a.s.l., 1–13 June 2022, det. Andreas Sanchez, specimen at MCSN.
- 2 females, Solorónch (Capriasca), Ticino, LV95: 2717643 E, 1101393 N (46.0538°N, 8.9587°E), 620 m a.s.l., 11 July–15 August 2022, det. Andreas Sanchez, specimens at MCSN.
- 54 females, Bolette (Locarno), Ticino, LV95: 2709912 E, 1112488 N (46.1549°N, 8.8614°E), 195 m a.s.l., 17 May–13 August 2022, det. Andreas Sanchez, specimens at MCSN and at Andreas Sanchez collection, Sion, Switzerland.
- 31 females, Bolette (Locarno), Ticino, LV95: 2709941 E, 1112476 N (46.1547°N, 8.8618°E), 195 m a.s.l., 2 June–13 August 2022, det. Andreas Sanchez, specimens at MCSN and at Andreas Sanchez collection, Sion, Switzerland.
- 3 females, Colombera (Stabio), Ticino, LV95: 2717680 E, 1078726 N (45.8499°N, 8.9535°E), 339 m a.s.l., 1–27 June 2022, det. Andreas Sanchez, specimens at MCSN.
- 5 females, Ronco del Re (Terre di Pedemonte), Ticino, LV95: 2699033 E, 1115853 N (46.1868°N, 8.7213°E), 372 m a.s.l., 17 May–23 August 2022, det. Andreas Sanchez, specimens at MCSN.
- 9 females, S. Martino (Vezia), Ticino, LV95: 2716328 E, 1098298 N (46.0262°N, 8.9409°E), 431 m a.s.l., 13 June–12 August 2022, det. Andreas Sanchez, specimens at MCSN and at Heiko Gebhardt collection, Tübingen, Germany.

Comparison of trap effectiveness

It was not an objective of the surveys reported here to compare the effectiveness of different trap types in capturing *A. maiche*. Nevertheless, a comparison of *A. maiche* captures per trap per day between trap types showed that bottle traps with ethanol as lure were more effective than green or black funnel traps with ethanol and additional attractants (Table 1). Captures of *A. maiche* during the main flight period (from June to August, see Fig. 3) at sites where the species occurred, averaged 1.02 ± 0.31 specimens per trap per day (mean ± SE, n = 11) for bottle traps with ethanol as lure (BT/E, trap type 1), 0.048 ± 0.018 specimens per trap per day (mean ± SE, n = 3) for green funnel traps with ethanol, alpha-pinene and eight-component blend (FG/EA8, trap type 2), 0.012 ± 0.007 specimens per trap per day (mean ± SE, n = 3) for black funnel traps with ethanol, alpha-pinene and eight-component blend (FB/EA8, trap type 3), and 0.073 ± 0.022 specimens per trap per day (mean ± SE, n = 29) for Polytraps with no lure (PT, trap type 4).

Other collection methods

No *A. maiche* were found infesting log sections (ca. 50 cm long, baited with ethanol) of European beech, Norway spruce and sweet chestnut that had been suspended from branches at study sites (1)–(6) in Ticino. However, five specimens were collected from around the cork used to



Figure 3. Mean trap captures of *Anisandrus maiche* in central-upper Ticino (left, Riviera-Iragna and Serravalle-Leggiuna, N = 4 ethanol-baited bottle traps) and in central Ticino (right, Bolette, Locarno, N = 2 Polytraps).

seal the ethanol reservoir of beech logs and one spruce log. No colonisation by *A. maiche* occurred of short sections of beech branches (about 20 cm long, 2–4 cm diameter, soaked in 70% ethanol) that had been placed on the ground at the same locations.

Phenology of *A. maiche* in Ticino

Captures of *A. maiche* with ethanol-baited bottle traps in central-upper Ticino (Riviera-Iragna and Serravalle-Leggiuna) (n = 4 traps) revealed that the main flight period was from June (or late May) to August (Fig. 3). This general flight pattern was observed also with the captures with unbaited Polytraps which occurred from mid-May to mid-August (Fig. 3) near Locarno at locations that were mainly at slightly lower elevations than the bottle trap sites in central Ticino. As the Polytraps were used from 10 March (i.e., late winter/early spring) until 3 October (i.e., early autumn), this indicates that no flights were missed, and that there is no second flight period. Therefore, the species is probably univoltine in the study region.

Fungal associates of *A. maiche*

From seven specimens of *A. maiche* that were caught alive, nine fungal cultures were successfully recovered. DNA barcoding confirmed that these cultures belonged to four species. Four cultures of *Ambrosiella cleistominuta* (including WSL DNA-ID PHP22_0914) (Ascomycota, Ceratocystidaceae) were obtained from sites Riviera-Iragna 2 and Serravalle-Leggiuna 1 (Fig. 4). In addition, *Aureobasidium pullulans* (de Bary & Löwenthal) G. Arnaud (Ascomycota, Dothioraceae; three cultures), *Cladosporium cladosporioides* (Fresen.) G.A. de Vries (Ascomycota, Davidiellaceae; one culture), and *Fusarium lateritium* Nees (Ascomycota, Nectriaceae; one culture) were found at Riviera-Iragna and Serravalle-Leggiuna sites. *Cladosporium cladosporioides* was isolated from the beetle surface, whereas the other three species were isolated from crushed beetles.

Figure 4. Culture of *Ambrosiella cleistominuta* (on 15-day old Potato Dextrose Agar, see methods) from *Anisandrus maiche* trapped alive in canton Ticino, Switzerland.

Discussion

Establishment and distribution of *A. maiche* in Switzerland

The detection of over 700 specimens of *Anisandrus maiche* across much of canton Ticino and an adjacent area of canton Grisons revealed that this species is well-established and already common in parts of southern Switzerland. This population appears to be contiguous with a recently detected population in adjacent parts of the Italian region of Lom-

bardy where specimens of *A. maiche* were found about 30 km from the nearest known occurrence in Switzerland (Ruzzier et al. 2022). In northern Italy, *A. maiche* was found in five provinces between Milan and Treviso and appears to be relatively widespread even though the first detection of this species occurred only in 2021 (Colombari et al. 2022; Ruzzier et al. 2022). It is very likely that *A. maiche* has been present in northern Italy and southern Switzerland already for many years. Its superficial resemblance to the common and widespread *Xylosandrus germanus* (Rabaglia et al. 2009), another non-native invasive ambrosia beetle of east Asian origin, probably prevented its earlier detection.

There is no indication that *A. maiche* occurs north of the Alpine divide (i.e., north or northwest of canton Ticino), although sampling and specific surveys for *A. maiche* north of the Alpine divide so far have been carried out only in parts of cantons Valais and Zurich. However, given the abundance of *A. maiche* in Ticino and northern Italy and the considerable volume of international and domestic trade, it is probably only a matter of time until *A. maiche* is established north of the Alpine divide. Most captures occurred at lower elevations in southern Ticino but some individuals were trapped at higher elevations (up to 915 m a.s.l.), indicating that mountain forests are also at risk of invasion by *A. maiche*. Previous studies have found a positive effect of ambient temperature on the number and activity density of non-native ambrosia beetles along with an effect of forest type (Rassati et al. 2016). However, it is likely that these patterns are also influenced by the locations where non-native species arrive, which tend to be urban and commercial areas in mainly warmer locations from where they disperse gradually.

Given that mountain forest habitats are particularly threatened by climate change (Engler et al. 2011), additional impacts by invasive species are a further concern. Although its climatic requirements have not been determined thoroughly, the presence of *A. maiche* in Ukraine (Terekhova and Skrylnik 2012), western Russia near Moscow (Nikitskii 2009) and in far-eastern Russian Siberia (Kurentsov 1941) suggests that the climatic conditions north of the Alps and at higher elevations in the mountains should be climatically suitable. However, species distribution modelling would be required to ascertain this with more certainty.

Effectiveness of different traps and lures

At sites where the species occurred, bottle traps baited with ethanol captured about 15 to 100 times more *A. maiche* individuals on average than funnel traps baited with ethanol and alpha-pinene or ethanol, alpha-pinene and an eight-component blend, or unbaited Polytrap flight interception traps. Previous studies have shown that ethanol by itself is a better lure for species of *Anisandrus* than a combination of ethanol and alpha-pinene, but this does not apply to all ambrosia beetles since species of *Xyleborus* and *Xylosandrus* showed different responses (Rabaglia et al. 2019). Catch rates can also be influenced by the concentration or release rate of ethanol (e.g., Cavaletto et al. 2021; Tobin and Ginzel 2023); however, this has not been studied specifically for *A. maiche*.

The apparent weakness of attraction in our experiments with ethanol-infused branch sections and log sections with an ethanol-reservoir was surprising given that others found similar methods to be effective for *A. maiche* (Reding and Ranger 2020) and other ambrosia beetle species (Monterrosa et al. 2021). It is possible that in our branch and log experiments, the attraction of *A. maiche* was compromised by a sub-optimal concentration and/or release rate of ethanol.

Trap colour and position (i.e., the height above ground level of traps) may also affect capture rates (e.g., Kerr et al. 2017; Cavaletto et al. 2020; Marchioro et al. 2020), although this has not been studied systematically for *A. maiche*. However, the fact that clear bottle traps baited with ethanol were highly attractive in our study suggests that trap colour appears not to be an important factor for *A. maiche*.

Phenology and host plants of *A. maiche*

Captures of *A. maiche* in Ticino occurred between early May and late August 2022. Although this stretches across a period of nearly four months, there was no clear indication of two separate peaks of flight activity, and we assume that the species is univoltine. Observations in Ukraine, Skrylnik et al. (2019) also suggest that *A. maiche* has one generation per year.

To our knowledge, only three infestations by *A. maiche* of trees are known from Ticino. Two were detected in 2023 at the botanic garden on the larger Brissago Island in Lago Maggiore. One concerned a live *Hakea* sp. shrub (Proteaceae), and the other a dead Chinese lacquer tree, *Toxicodendron vernicifluum* (Anacardiaceae). The third infestation was in a twig (5 mm diameter) of a chestnut tree at a forest edge east of Tegna, Ticino. Considering its wide distribution and apparent abundance in Ticino, the lack of any other observed infestations is surprising. However, based on host records from its native and non-native ranges (summarised in Hölling and Brockerhoff 2023; Ruzzier et al. 2023), *A. maiche* is known to be highly polyphagous. In its native range in eastern Asia, host records include trees in the genera *Acer* (maple), *Alnus* (alder), *Betula* (birch), *Carpinus* (hornbeam), *Corylus* (hazel), *Fraxinus* (ash), *Juglans* (walnut), *Quercus* (oak), *Ulmus* (elm), and several other genera, including some conifers (Hölling and Brockerhoff 2023). In the non-native range of *A. maiche*, relatively few host records exist which, nonetheless, confirm that this species has a very wide host range. In Ukraine, attacks of *Betula pendula*, *Populus tremula*, *Quercus robur*, *Quercus rubra* (a North American oak species) and *Ulmus minor* have been reported (Nikulina et al. 2015; Skrylnik et al. 2019). In North America, *A. maiche* has been observed breeding in *Acer rubrum*, *Cercis canadensis*, *Cornus florida*, *Gleditsia triacanthos*, *Quercus imbricaria*, *Sassafras albidum* and *Styrax japonicus* (Mayers et al. 2017; Ranger et al. 2015, 2019, 2020; Reding and Ranger 2020).

Introduction pathways for *A. maiche* and other ambrosia beetles

To our knowledge, no confirmed border interceptions of *A. maiche* with any traded goods have been recorded, neither in Switzerland, anywhere else within the EPPO region, nor in a number of other countries (Brockerhoff et al. 2006; Turner et al. 2021). However, many other ambrosia beetles including species in the genera *Xyleborus*, *Xyleborinus*, *Xylosandrus* and others have been intercepted numerous times (Brockerhoff et al. 2006; Haack et al. 2006, 2013), indicating the existence of pathways that can facilitate invasions of ambrosia beetles. For example, *Xylosandrus crassiusculus* and *Xylosandrus germanus*, two well-known invasive ambrosia beetles that are now established in several continents where they are not native, have been intercepted repeatedly in New Zealand (Brockerhoff et al. 2006). Furthermore, it is possible that specimens of *A. maiche* were intercepted in some countries but not recognised as this species. Pathways known or thought to be involved in invasions of ambrosia beetles include international transport of wood packaging materials (such as pallets and case wood used with ceramic tiles, stone products and numerous other commodities) as well as trade in firewood and live plants (Liebhold et al. 2012; Meurisse et al. 2019). According to Mandelshtam et al. (2018), it may be possible that *A. maiche* extended its distribution by natural spread westwards from its native range in the Russian Far East; however, considering its apparently discontinuous distribution between eastern Asia and central Europe, we regard it as more likely that international trade of infested wood or wood products is responsible for its arrival and establishment in Europe.

Fungal associates of *A. maiche*

Ambrosiella cleistominuta is the ambrosia fungus associated with *Anisandrus maiche* (Mayers et al. 2017). Previously, this association was only known from the United States, where the fungus was isolated and described as a species new to science in 2017 (Mayers et al. 2017). Here, we report for the first time that the fungus is also present in Europe, again in association with *A. maiche* (with four confirmed isolations from *A. maiche* in the present study). The fungus was most likely introduced to Europe together with its beetle host. We are not aware of any studies indicating that this fungus is pathogenic to (live) plants.

Surprisingly, *A. cleistominuta* was also found in association with *Xylosandrus crassiusculus* (Motschulsky 1866) in samples of the same *Hakea* shrub from the larger Brissago Island which was also infested by *A. maiche*. This was first reported by Ribeiro-Correia et al. (2023), and a more detailed publication on this case is in preparation. *Xylosandrus crassiusculus* is a recent invader to Switzerland which was first detected in Ticino in 2013 (Sanchez et al. 2020). *Xylosandrus crassiusculus* is usually always associated with a different ambrosia fungus, *Ambrosiella roeperi* T.C. Harr. & McNew (Harrington et al. 2014; Saragih et al. 2021). Ambrosia beetles and their mutualistic ambrosia fungi typically show a high degree of fidelity, and associations with other ambrosia fungi are rare (Biedermann and Vega 2020). However, new associations between ambrosia beetles and ambrosia fungi can be established experimentally (Menocal et al. 2023) and may thus also occur in nature, in particular as a result of invasions of ambrosia beetles into non-native regions. "Adoptions" of foreign ambrosia fungi of beetles colonizing the same trees in the invasive range may be sufficient for nutrition and development in the foreign gallery, but may not be suitable for transport to other trees because of the evolved specificity of mycangia (Mayers et al. 2022). Successful mycangial transmission should only occur in beetle species with similar mycangial selection processes and that are obligately associated with species in the same fungal clade (Biedermann and Vega 2020). The genus *Ambrosiella* (Ascomycota: Microascales), for example, is mutualistic with scolytine ambrosia beetles in the *Xylosandrus* and *Anisandrus* genera, so fungal exchanges between these beetle species are most likely (Mayers et al. 2015). Newly acquired ambrosia fungi may then lead to reductions or even increases in beetle fitness (e.g., reproductive success) (Menocal et al. 2023).

Three other fungi were isolated from *A. maiche* trapped alive in Ticino: *Aureobasidium pullulans* is a saprophytic yeast-like fungus with a worldwide distribution; it occurs on the leaves of a wide range of plants and is known mainly from crop plants (Deshpande et al. 1992). *Cladosporium cladosporioides* is a member of a genus that includes the most common environmental fungi found worldwide with species of various lifestyles (Bensch et al. 2012). *Cladosporium cladosporioides* is also recognized as a pathogen of grapes and other fruits on which it produces the so-called *Cladosporium* fruit rot (Latorre et al. 2011; Swett et al. 2019). Its isolation from the body surface of *A. maiche* may suggest a coincidental association of the fungus and the beetle. *Fusarium lateritium* is a globally distributed plant pathogen that may cause a variety of symptoms on affected plants. For example, *F. lateritium* can cause chlorotic leaf distortion on sweet potato (Clark et al. 1995) and nut grey necrosis on hazelnut fruits (Vitale et al. 2011). Recently, *F. lateritium* was found to cause shoot dieback of boxelder maple (*Acer negundo*) in Poland (Patejuk et al. 2022). It is not yet known whether any of these three fungi (and their association with *A. maiche*) could pose a problem for forest trees in Switzerland and Central Europe. As they have also only been found sporadically with *A. maiche*, we suspect that these may be coincidental associations. Nevertheless, these examples show that *A. maiche* can be a potential vector of pathogens.

Potential damage to infested trees from *A. maiche* and associated fungi

So far, the only trees (or wood) infested by *A. maiche* that have been found in Switzerland (the vast majority of specimens were trapped) are two non-native plant species at a botanic garden, a *Hakea* sp. shrub (Proteaceae), and a

dead Chinese lacquer tree, *Toxicodendron vernicifluum* (Anacardiaceae) and a native chestnut tree at the edge of a semi-natural forest. It is uncertain what other tree or shrub species are attacked in Switzerland. However, most of the trap catches of *A. maiche* in canton Ticino occurred at locations with native forest and native tree species. Based on this as well as previous host records (summarised in Hölling and Brockerhoff (2023) and Ruzzier et al. (2023)), *A. maiche* may attack a wide range of broadleaved tree species both native and non-native present in Ticino.

In general, most ambrosia beetles attack freshly dead wood and thus do not cause much damage of live plants, but there are exceptions, particularly when non-native ambrosia beetles and associated plant pathogenic fungi are involved (Hulcr and Dunn 2011; Eskalen et al. 2013; Hughes et al. 2017; Paap et al. 2018; Morales-Rodríguez et al. 2021). Based on observations from Ukraine, Terekhova and Skrylnik (2012) state that *A. maiche* "has no significant economic impact" in that country. However, "noticeable" damage to birch (*Betula pendula*) trees in Ukraine was described by Skrylnik et al. (2019) who gave *A. maiche* an intermediate impact rating for birch with a "physiological harmfulness score" of 5 out of 14, considering both damage from galleries made by the beetles and their assumed ability to act as vectors of plant pathogens. As of late 2023, there is no indication of any major damage caused by *A. maiche* in Switzerland, neither from its galleries nor from any associated plant pathogens. However, no systematic surveys for damage from *A. maiche* have been carried out yet, and it is not known if *A. maiche* is indeed harmful to trees or shrubs. Nevertheless, its considerable abundance in some locations in Ticino along with the apparent association with at least one plant pathogen (*F. lateritium*, see Patejuk et al. (2022)) suggests that there is some potential for damage, especially if plants are stressed by drought, flooding or fire (Terekhova and Skrylnik 2012; Ranger et al. 2015; Mandelshtam et al. 2018). Ranger et al. (2015) consider that *A. maiche* prefers to attack living, weakened trees over dead trees.

Further research needs

Several aspects of the invasion of *A. maiche* in Europe deserve further study. As this species is already relatively common in southern Switzerland (and probably also in other invaded areas), it is possible that it will become one of the most abundant ambrosia beetles. This has happened at some locations in North America, where *A. maiche* was found to be one of the two most abundant ambrosia beetles, together with *Xylosandrus germanus* (Ranger et al. 2019). An increase in the abundance of *A. maiche* could lead to more noticeable damage of infested trees (given its apparent preference for attacking weakened living trees) and it could also cause competition with native ambrosia beetles which may decline in response. Therefore, having a better understanding of the extent of the distribution of *A. maiche* both at higher elevations in mountain forests of Ticino and beyond canton Ticino is desirable. Likewise, better knowledge of its host plants, fungal associates and their combined potential effect on plant health are required in both lowland and mountain forests. Furthermore, a better understanding of the pathways involved in the invasions and spread of *A. maiche* and other ambrosia beetles could be used to limit the extent of future invasions and impacts. Finally, given the occurrence of heteroplasmy in COI sequences in some of the samples of *A. maiche* we analysed, a known problem with COI barcoding of insects (e.g., Magnacca and Brown 2010; Cognato et al. 2020), further studies on the invasion and population genetics of *A. maiche* should examine the use of additional markers.

Acknowledgements

We thank Adrian Oncelli (Repubblica e Cantone Ticino, Dipartimento del territorio, Sezione forestale), Simone Serretti (Repubblica e Cantone Ticino, Dipartimento del territorio, Ufficio della natura e del paesaggio), Bärbel Koch and Lucia Pollini Paltrinieri from the Natural History Museum of the Canton Ticino for their support and contributions to trapping, the regional forestry offices for their help, namely Cristian Gobbin, Aron Ghiringhelli, Daniele Barra, Nicola Bomio-Pacciorini, Jacques Bottani, Patrick Luraschi, Thomas Schiesser, Giovanni Galli, Martino Bonardi, and the landowners and citizens communities (patriziati) for their permission to sample on their land, Aline Knoblauch (Federal Office for the Environment (FOEN), Forest Division) for project support, Vienna Kowallik (Universität Freiburg, Forstentomologie und Waldschutz) for training in fungal extraction, Bjarte Jordal (University of Bergen, Natural History Collections) for sharing a specimen, Quirin Kupper, Sven Ulrich und Robin Winiger (WSL) for laboratory support, Carolina Cornejo (WSL) for interpretation of DNA sequences, and Carl-Michael Anderson (WSL) for taking photos of *A. maiche*. We further thank Stefano Brignoli, Ivan Candolfi, Lisa Elzi, Giorgio Musso and Davide Prati for their work in the field and insect sorting in the lab. This project was supported in part by funding from the Swiss Federal Office for the Environment (FOEN/OFEV/BAFU) to WSL and the Canton of Ticino. MK was partly supported by the Ministry of Agriculture of the Czech Republic, institutional support MZE-RO0123.

References

Alonso-Zarazaga MA, Barrios H, Borovec R, Bouchard P, Caldara R, Colonnelli E, Gültekin L, Hlaváč P, Korotyaev B, Lyal CHC, Machado A, Meregalli M, Pierotti H, Ren L, Sánchez-Ruiz M, Sforzi A, Silfverberg H, Skuhrovec J, Trýzna M, Velázquez de Castro AJ, Yunakov NN (2017) Cooperative Catalogue of Palaearctic Coleoptera Curculionoidea. Monografías electrónicas SEA, Vol. 8. Sociedad Entomológica Aragonesa S.E.A., Zaragoza, 729 pp. https://www.biotaxa.org/mesea/article/view/34195/30253

Alonso-Zarazaga MA, Barrios H, Borovec R, Bouchard P, Caldara R, Colonnelli E, Gültekin L, Hlaváč P, Korotyaev B, Lyal CHC, Mach-

ado A, Meregalli M, Pierotti H, Ren L, Sánchez-Ruiz M, Sforzi A, Silfverberg H, Skuhrovec J, Trýzna M, Velázquez de Castro AJ, Yunakov NN (2023) Cooperative Catalogue of Palaearctic Coleoptera Curculionoidea (2nd edn.). Monografías electrónicas SEA (Vol. 14). Sociedad Entomológica Aragonesa S.E.A., Zaragoza, 780 pp.

Bensch K, Braun U, Groenewald JZ, Crous PW (2012) The genus *Cladosporium*. Studies in Mycology 72(1): 1–401. https://doi.org/10.3114/sim0003

Biedermann PH, Vega FE (2020) Ecology and evolution of insect–fungus mutualisms. Annual Review of Entomology 65: 431–455. https://doi.org/10.1146/annurev-ento-011019-024910

Brockerhoff EG, Bain J, Kimberley M, Knížek M (2006) Interception frequency of exotic bark and ambrosia beetles (Coleoptera: Scolytinae) and relationship with establishment in New Zealand and worldwide. Canadian Journal of Forest Research 36(2): 289–298. https://doi.org/10.1139/x05-250

Brockerhoff EG, Liebhold AM (2017) Ecology of forest insect invasions. Biological Invasions 19: 3141–3159. https://doi.org/10.1007/s10530-017-1514-1

Brustel H (2012) PolytrapTM 2010: new "soft design" window flight trap for saproxylic beetles. In: Jurc M (Ed.) Saproxylic Beetles in Europe: Monitoring, Biology and Conservation. Slovenian Forestry Institute, Silva Slovenica, 91–92.

Cavaletto G, Faccoli M, Marini L, Spaethe J, Magnani G, Rassati D (2020) Effect of trap color on captures of bark-and wood-boring beetles (Coleoptera; Buprestidae and Scolytinae) and associated predators. Insects 11: e749. https://doi.org/10.3390/insects11110749

Cavaletto G, Faccoli M, Ranger CM, Rassati D (2021) Ambrosia beetle response to ethanol concentration and host tree species. Journal of Applied Entomology 145: 800–809. https://doi.org/10.1111/jen.12895

Clark CA, Hoy MW, Nelson PE (1995) Variation among isolates of *Fusarium lateritium* from sweetpotato for pathogenicity and vegetative compatibility. Phytopathology 85: 624–629. https://doi.org/10.1094/Phyto-85-624

Colombari F, Martinez-Sañudo I, Battisti A (2022) First report of the alien ambrosia beetle *Cnestus mutilatus* and further finding of *Anisandrus maiche* in the European part of the EPPO region (Coleoptera: Curculionidae: Scolytinae: Xyleborini). EPPO Bulletin 52: 446–450. https://doi.org/10.1111/epp.12840

Cognato AI, Sari G, Smith SM, Beaver RA, Li Y, Hulcr J, Jordal BH, Kajimura H, Lin CS, Pham TH, Singh S (2020) The essential role of taxonomic expertise in the creation of DNA databases for the identification and delimitation of Southeast Asian ambrosia beetle species (Curculionidae: Scolytinae: Xyleborini). Frontiers in Ecology and Evolution 8: 1–27. https://doi.org/10.3389/fevo.2020.00027

Deshpande MS, Rale VB, Lynch JM (1992) *Aureobasidium pullulans* in applied microbiology: a status report. Enzyme and Microbial Technology 14(7): 514–527. https://doi.org/10.1016/0141-0229(92)90122-5

Dole SA, Jordal BH, Cognato AI (2010) Polyphyly of *Xylosandrus* Reitter inferred from nuclear and mitochondrial genes (Coleoptera: Curculionidae: Scolytinae). Molecular Phylogenetics and Evolution 54(3): 773–782. https://doi.org/10.1016/j.ympev.2009.11.011

Engler R, Randin CF, Thuiller W, Dullinger S, Zimmermann NE, Araujo MB, Pearman PB, Le Lay G, Piedallu C, Albert CH, Choler P (2011) 21st century climate change threatens mountain flora unequally across Europe. Global Change Biology 17(7): 2330–2341. https://doi.org/10.1111/j.1365-2486.2010.02393.x

EPPO (2022) EPPO Global Database – *Anisandrus maiche* (ANIDMA). https://gd.eppo.int/taxon/ANIDMA/hosts

Eskalen A, Stouthamer R, Lynch SC, Rugman-Jones PF, Twizeyimana M, Gonzalez A, Thibault T (2013) Host range of *Fusarium* dieback and its ambrosia beetle (Coleoptera: Scolytinae) vector in southern California. Plant Disease 97: 938–951. https://doi.org/10.1094/PDIS-11-12-1026-RE

Fan JT, Denux O, Courtin C, Bernard A, Javal M, Millar JG (2019) Multi-component blends for trapping native and exotic longhorn beetles at potential points-of-entry and in forests. Journal of Pest Science 92: 281–297. https://doi.org/10.1007/s10340-018-0997-6

Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3(5): 294–299.

Fraedrich SW, Harrington TC, Rabaglia RJ, Ulyshen MD, Mayfield Iii AE, Hanula JL, Eickwort JM, Miller DR (2008) A fungal symbiont of the redbay ambrosia beetle causes a lethal wilt in redbay and other Lauraceae in the southeastern United States. Plant Disease 92: 215–224. https://doi.org/10.1094/PDIS-92-2-0215

Franić I, Prospero S, Hartmann M, Allan E, Auger-Rozenberg M-A, Grünwald N, Kenis M, Roques A, Schneider S, Sniezko R, Williams W, Eschen R (2019) Are traded forest tree seeds a potential source for alien pests? Ecological Applications 29(7): e01971. https://doi.org/10.1002/eap.1971

Gomez DF, Rabaglia RJ, Fairbanks KEO, Hulcr J (2018) North American Xyleborini north of Mexico: a review and key to genera and species (Coleoptera, Curculionidae, Scolytinae). ZooKeys 768: 19–68. https://doi.org/10.3897/zookeys.768.24697

Gossner MM, Falck K, Weisser WW (2019) Effects of management on ambrosia beetles and their antagonists in European beech forests. Forest Ecology and Management 437: 126–133. https://doi.org/10.1016/j.foreco.2019.01.034

Grégoire JC, Piel F, De Proft M, Gilbert M (2001) Spatial distribution of ambrosia-beetle catches: a possibly useful knowledge to improve mass-trapping. Integrated Pest Management Reviews 6: 237–242. https://doi.org/10.1023/A:1025723402355

Grüne S (1979) Handbuch zur Bestimmung der Europäischen Borkenkäfer (Brief illustrated key to European bark beetles). Verlag M & H Schapfer, Hanover, 182 pp.

Haack RA (2006) Exotic bark-and wood-boring Coleoptera in the United States: Recent establishments and interceptions. Canadian Journal of Forest Research 36(2): 269–288. https://doi.org/10.1139/x05-249

Haack RA, Rabaglia RJ (2013) Exotic bark and ambrosia beetles in the USA: potential and current invaders. In: Peña J (Ed.) Potential Invasive Pests of Agricultural Crops. CABI: 48–74. https://doi.org/10.1079/9781845938291.0048

Harrington TC, Fraedrich SW, Aghayeva DN (2008) *Raffaelea lauricola*, a new ambrosia beetle symbiont and pathogen on the Lauraceae. Mycotaxon 104: 399–404.

Harrington TC, McNew D, Mayers C, Fraedrich SW, Reed SE (2014) *Ambrosiella roeperi* sp. nov. is the mycangial symbiont of the granulate ambrosia beetle, *Xylosandrus crassiusculus*. Mycologia 106(4): 835–845. https://doi.org/10.3852/13-354

Hölling D, Brockerhoff EG (2023) Express Pest Risk Analysis (PRA) for: *Anisandrus maiche* (Kurentzov). Prepared by the Swiss Federal Institute for Forest, Snow and Landscape Research for the Swiss Federal Office for the Environment. Bern.

Hughes MA, Riggins JJ, Koch FH, Cognato AI, Anderson C, Formby JP, Dreaden TJ, Ploetz RC, Smith JA (2017) No rest for the laurels: symbiotic invaders cause unprecedented damage to southern USA forests. Biological Invasions 19: 2143–2157. https://doi.org/10.1007/s10530-017-1427-z

Hulcr J, Dunn RR (2011) The sudden emergence of pathogenicity in insect-fungus symbioses threatens native forest ecosytems. Proceedings of the Royal Society B: Biological Sciences 278: 2866–2873. https://doi.org/10.1098/rspb.2011.1130

Hulcr J, Stelinski LL (2017) The ambrosia symbiosis: from evolutionary ecology to practical management. Annual Review of Entomology 62: 285–303. https://doi.org/10.1146/annurev-ento-031616-035105

Kambestad M, Kirkendall LR, Knutsen IL, Jordal BH (2017) Cryptic and pseudo-cryptic diversity in the world's most common bark beetle – *Hypothenemus eruditus*. Organisms Diversity and Evolution 17: 633–652. https://doi.org/10.1007/s13127-017-0334-6

Kenis M, Auger-Rozenberg M-A, Roques A, Timms L, Péré C, Cock MJW, Settele J, Augustin S, Lopez-Vaamonde C (2009) Ecological effects of invasive alien insects. Biological Invasions 11: 21–45. https://doi.org/10.1007/978-1-4020-9680-8_3

Kerr JL, Kelly D, Bader MK-F, Brockerhoff EG (2017) Olfactory cues, visual cues, and semiochemical diversity interact during host location by invasive forest beetles. Journal of Chemical Ecology 43: 17–25. https://doi.org/10.1007/s10886-016-0792-x

Kirkendall LR (1983) The evolution of mating systems in bark and ambrosia beetles (Coleoptera: Scolytidae and Platypodidae). Zoological Journal of the Linnean Society 77(4): 293–352. https://doi.org/10.1111/j.1096-3642.1983.tb00858.x

Kirkendall LR, Faccoli M (2010) Bark beetles and pinhole borers (Curculionidae, Scolytinae, Platypodinae) alien to Europe. ZooKeys 56: 227–251. https://doi.org/10.3897/zookeys.56.529

Knížek M (2011) Scolytinae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 7). Curculionoidea I. Apollo Books, Stenstrup, 204–251.

Kovalenko YN, Nikitskii NB (2013) New to Russia and interesting finds of xylophilous beetles (Coleoptera) in the middle Russian forest-steppe zone. Bjulleten' Moskovskogo Obscestva Ispytatelej Prirody / Otdel biologiceskij 118(2): 20–26.

Kurentsov AI (1941) Koroedy Dal'nego Vostoka SSSR (The Bark Beetles of Far East of USSR). Izdatl'stvo Akademii nauk SSSR, Moscow, 235 pp.

Lantschner MV, Corley JC, Liebhold AM (2020) Drivers of global Scolytinae invasion patterns. Ecological Applications 30(5): e02103. https://doi.org/10.1002/eap.2103

Latorre BA, Briceño EX, Torres R (2011) Increase in *Cladosporium* spp. populations androt of wine grapes associated with leaf removal. Crop Protection 30: 52–56. https://doi.org/10.1016/j.cropro.2010.08.022

Liebhold AM, Brockerhoff EG, Garrett LJ, Parke JL, Britton KO (2012) Live plant imports: the major pathway for forest insect and pathogen invasions of the US. Frontiers in Ecology and the Environment 10(3): 135–143. https://doi.org/10.1890/110198

Magnacca KN, Brown MJF (2010) Mitochondrial heteroplasmy and DNA barcoding in Hawaiian *Hylaeus* (*Nesoprosopis*) bees (Hymenoptera: Colletidae). BMC Evolutionary Biology 10: e174. https://doi.org/10.1186/1471-2148-10-174

Mandelshtam MY, Yakushkin EA, Petrov AV (2018) Oriental ambrosia beetles (Coleoptera: Curculionidae: Scolytinae): new inhabitants of Primorsky krai in Russia. Russian Journal of Biological Invasions 9: 355–365. https://doi.org/10.1134/S2075111718040082

Marchioro M, Rassati D, Faccoli M, Van Rooyen K, Kostanowicz C, Webster V, Mayo P, Sweeney J (2020) Maximizing bark and ambrosia beetle (Coleoptera: Curculionidae) catches in trapping surveys for longhorn and jewel beetles. Journal of Economic Entomology 113(6): 2745–2757. https://doi.org/10.1093/jee/toaa181

Mayers CG, McNew DL, Harrington TC, Roeper RA, Fraedrich SW, Biedermann PH, Castrillo LA, Reed SE (2015) Three genera in the Ceratocystidaceae are the respective symbionts of three independent lineages of ambrosia beetles with large, complex mycangia. Fungal Biology 119(11): 1075–1092. https://doi.org/10.1016/j.funbio.2015.08.002

Mayers CG, Harrington TC, Ranger CM (2017) First report of sexual state in an ambrosia fungus: *Ambrosiella cleistominuta* sp. nov. associated with the ambrosia beetle *Anisandrus maiche*. Botany 95: 503–512. https://doi.org/10.1139/cjb-2016-0297

Mayers CG, Harrington TC, Biedermann PHW (2022) Mycangia define the diverse ambrosia beetle–fungus symbioses. In: Schultz TR, Gawne R, Peregrine PN (Eds) The Convergent Evolution of Agriculture in Humans and Insects. The MIT Press, London, 105–142. https://doi.org/10.7551/mitpress/13600.003.0013

Menocal O, Cruz LF, Kendra PE, Berto M, Carrillo D (2023) Flexibility in the ambrosia symbiosis of *Xyleborus bispinatus*. Frontiers in Microbiology 14: e1110474. https://doi.org/10.3389/fmicb.2023.1110474

Meurisse N, Rassati D, Hurley BP, Brockerhoff EG, Haack RA (2019) Common pathways by which non-native forest insects move internationally and domestically. Journal of Pest Science 92: 13–27. https://doi.org/10.1007/s10340-018-0990-0

Monterrosa A, Acebes AL, Blaauw B, Joseph SV (2021) Effects of trap, and ethanol lure type and age on attraction of ambrosia beetles (Coleoptera: Curculionidae). Journal of Economic Entomology 114: 1647–1654. https://doi.org/10.1093/jee/toab089

Morales-Rodríguez C, Sferrazza I, Aleandri MP, Dalla Valle M, Speranza S, Contarini M, Vannini A (2021) The fungal community associated with the ambrosia beetle *Xylosandrus compactus* invading the Mediterranean maquis in central Italy reveals high biodiversity and suggests environmental acquisitions. Fungal Biology 125(1): 12–24. https://doi.org/10.1016/j.funbio.2020.09.008

Nikitskii NB (2009) New and interesting finds of xylophilic and some other darkling beetles (Coleoptera) in Moscow oblast and Moscow. Bjulleten' Moskovskogo Obscestva Ispytatelej Prirody / Otdel biologiceskij 114(5): 50–58. https://doi.org/10.23885/1814-3326-2013-9-2-223-241

Nikulina TV, Martynov VV, Mandel'shtam MY (2007) *Anisandrus maiche* – A new species of the bark beetles (Coleoptera, Scolytidae) in European fauna. Vestnik Zoologii 41(6): 1–542. https://doi.org/10.23885/1814-3326-2013-9-2-223-241

Nikulina T, Mandelshtam M, Petrow A, Nazarenko V, Yunakov N (2015) A survey of weevils of Ukraine. Bark and Ambrosia beetles (Coleoptera: Curculionidae: Platypodinae and Scolytinae). Zootaxa 3912: 1–61. https://doi.org/10.11646/zootaxa.3912.1

Paap T, de Beer ZW, Migliorini D, Nel WJ, Wingfield MJ (2018) The polyphagous shot hole borer (PSHB) and its fungal symbiont *Fusarium euwallaceae*: a new invasion in South Africa. Australasian Plant Pathology 47: 231–237. https://doi.org/10.1007/s13313-018-0545-0

Park S, Smith SM, Cognato AI, Beaver RA (2020) Catalogue of Korean Xyleborine ambrosia beetles (Coleoptera: Curculionidae: Scolytinae) with seven new species. Journal of Asia-Pacific Biodiversity 13(2): 210–228. https://doi.org/10.1016/j.japb.2020.01.002

Patejuk K, Baturo-Cieśniewska A, Najberek K, Pusz W (2022) First report of *Fusarium lateritium* causing shoot dieback of *Acer negundo* in Europe. Plant Disease 106(5): e1519. https://doi.org/10.1094/PDIS-06-21-1294-PDN

Pfeffer A (1995) Zentral und Westpaläarktische Borken und Kernkäfer. Naturhistorisches Museum, Basel, 310 pp.

PGesV (2018) Verordnung vom 31. Oktober 2018 über den Schutz von Pflanzen vor besonders gefährlichen Schadorganismen (Pflanzengesundheitsverordnung, PGesV; SR 916.20). Schweizerische Eidgenossenschaft. https://www.fedlex.admin.ch/eli/cc/2018/682/de

PGesV-WBF-UVEK (2019) Verordnung vom 14. November 2019 des WBF und des UVEK zur Pflanzengesundheitsverordnung (PGesV-WBF-UVEK; SR 916.201). Schweizerische Eidgenossenschaft. https://www.fedlex.admin.ch/eli/cc/2019/787/de

Pyšek P, Hulme PE, Simberloff D, Bacher S, Blackburn TM, Carlton JT, Dawson W, Essl F, Foxcroft LC, Genovesi P, Jeschke JM, Kühn I, Liebhold AM, Mandrak NE, Meyerson LA, Pauchard A, Pergl J, Roy HE, Seebens H, van Kleunen M, Vilà M, Wingfield MJ, Richardson DM (2020) Scientists' warning on invasive alien species. Biological Reviews 95(6): 1511–1534. https://doi.org/10.1111/brv.12627

Rabaglia RJ, Vandenberg NJ, Acciavatti RE (2009) First records of *Anisandrus maiche* Stark (Coleoptera: Curculionidae: Scolytinae) from North America. Zootaxa 2137: 23–28. https://doi.org/10.11646/zootaxa.2137.1.2

Rabaglia RJ, Cognato AI, Hoebeke ER, Johnson CW, LaBonte JR, Carter ME, Vlach JJ (2019) Early detection and rapid response: a 10-year summary of the USDA Forest Service program of surveillance for non-native bark and ambrosia beetles. American Entomologist 65(1): 29–42. https://doi.org/10.1093/ae/tmz015

Ranger CM, Schultz PB, Frank SD, Chong JH, Reding ME (2015) Non-native ambrosia beetles as opportunistic exploiters of living but weakened trees. PLOS ONE 10: e0131496. https://doi.org/10.1371/journal.pone.0131496

Ranger CM, Reding ME, Schultz PB, Oliver JB, Frank SD, Addesso KM, Chong JH, Sampson B, Werle C, Gill S, Krause C (2016) Biology, ecology, and management of nonnative ambrosia beetles (Coleoptera: Curculionidae: Scolytinae) in ornamental plant nurseries. Journal of Integrated Pest Management 7(1): 1–9. https://doi.org/10.1093/jipm/pmw005

Ranger CM, Schultz PB, Frank SD, Reding ME (2019) Freeze stress of deciduous trees induces attacks by opportunistic ambrosia beetles. Agricultural and Forest Entomology 21: 168–179. https://doi.org/10.1111/afe.12317

Ranger CM, Werle CT, Schultz PB, Addesso KM, Oliver JB, Reding ME (2020) Long-lasting insecticide netting for protecting tree stems from attack by ambrosia beetles (Coleoptera: Curculionidae: Scolytinae). Insects 11: 1–8. https://doi.org/10.3390/insects11010008

Rassati D, Faccoli M, Battisti A, Marini L (2016) Habitat and climatic preferences drive invasions of non-native ambrosia beetles in deciduous temperate forests. Biological Invasions 18: 2809–2821. https://doi.org/10.1007/s10530-016-1172-8

Reding ME, Ranger CM (2020) Attraction of invasive ambrosia beetles (Coleoptera: Curculionidae: Scolytinae) to ethanol-treated tree bolts. Journal of Economic Entomology 113: 321–329. https://doi.org/10.1093/jee/toz282

Ribeiro-Correia JP, Prospero S, Beenken L, Biedermann PH, Blaser S, Chittaro Y, Frey D, Hölling D, Kaya SO, Knížek M, Mittelstrass J, Branco M, Ruffner B, Sanchez A, Brockerhoff EG (2023) Distribution of the invasive *Anisandrus maiche* (Coleoptera: Scolytinae) in Switzerland, first record in Europe of its ambrosia fungus *Ambrosiella cleistominuta*, and its new association with *Xylosandrus crassiusculus*. bioRxiv: 2023.03.30.534995. https://doi.org/10.1101/2023.03.30.534995

Roques A, Ren L, Rassati D, Shi J, Akulov E, Audsley N, Auger-Rozenberg M-A, Avtzis D, Battisti A, Bellanger R, Bernard A, Bernadinelli I, Branco M, Cavaletto G, Cocquempot C, Contarini M, Courtial B, Courtin C, Denux O, Dvořák M, Fan J-t, Feddern N, Francese J, Franzen EKL, Garcia A, Georgiev G, Georgieva M, Giarruzzo F, Gossner M, Gross L, Guarneri D, Hoch G, Hölling D, Jonsell M, Kirichenko N, Loomans A, Luo Y-q, McCullough D, Maddox C, Magnoux E, Marchioro M, Martinek P, Mas H, Mériguet B, Pan Y-z, Phélut R, Pineau P, Ray AM, Roques O, Ruiz M-C, Sarto i Monteys V, Speranza S, Sun J-h, Sweeney JD, Touroult J, Valladares L, Veillat L, Yuan Y, Zalucki MP, Zou Y, Žunič-Kosi A, Hanks LM, Millar JG (2023) Worldwide tests of generic attractants, a promising tool for early detection of non-native cerambycid species. NeoBiota 84: 169–209. https://doi.org/10.3897/neobiota.84.91096

Ruzzier E, Bani L, Cavaletto G, Faccoli M, Rassati D (2022) *Anisandrus maiche* Kurentzov (Curculionidae: Scolytinae), an Asian species recently introduced and now widely established in Northern Italy. BioInvasions Records 11(3): 652–658. https://doi.org/10.3391/bir.2022.11.3.07

Ruzzier E, Ortis G, Vallotto D, Faccoli M, Martinez-Sañudo I, Marchioro M (2023) The first full host plant dataset of Curculionidae Scolytinae of the world: tribe Xyleborini LeConte, 1876. Scientific Data 10(1): e166. https://doi.org/10.1038/s41597-023-02083-5

Sanchez A, Chittaro Y, Germann C, Knížek M (2020) Annotated checklist of Scolytinae and Platypodinae (Coleoptera, Curculionidae) of Switzerland. Alpine Entomology 4: 81–97. https://doi.org/10.3897/alpento.4.50440

Sanchez A, Chittaro Y, Frey D, Koch B, Hölling D, Brockerhoff EG, Ruffner B, Knížek M (2023) Three alien ambrosia beetles (Coleoptera: Curculionidae, Scolytinae) new to Switzerland. Alpine Entomology 7: 45–55. https://doi.org/10.3897/alpento.7.103269

Saragih SA, Takemoto S, Kusumoto D, Kamata N (2021) Fungal diversity in the mycangium of an ambrosia beetle *Xylosandrus crassiusculus* (Coleoptera: Curculionidae) in Japan during their late dispersal season. Symbiosis 84: 111–118. https://doi.org/10.1007/s13199-021-00762-8

Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, et al. (2018) Global rise in emerging alien species results from increased accessibility of new source pools. Proceedings of the National Academy of Sciences 115: E2264–E2273. https://doi.org/10.1073/pnas.1719429115

Skrylnik Y, Koshelyaeva Y, Meshkova V (2019) Harmfulness of xylophagous insects for silver birch (*Betula pendula* Roth.) in the left-bank forest-steppe of Ukraine. Folia Forestalia Polonica, Series A – Forestry 61: 159–173. https://doi.org/10.2478/ffp-2019-0016

Smith SM, Beaver RA, Cognato AI (2020) A monograph of the Xyleborini (Coleoptera, Curculionidae, Scolytinae) of the Indochinese Peninsula (except Malaysia) and China. ZooKeys 983: 1–442. https://doi.org/10.3897/zookeys.983.52630

Stark VN (1952) Zhestkokrylye, Koroedy. Fauna SSSR (Vol. 31). Akademia Nauk SSSR: 1–463.

Swett CL, Hamby KA, Hellman EM, Carignan C, Bourret TB, Koivunen EE (2019) Characterizing members of the *Cladosporium cladosporioides* species complex as fruit rot pathogens of red raspberries in the mid-Atlantic and co-occurrence with *Drosophila suzukii* (spot-

ted wing drosophila). Phytoparasitica 47: 415–428. https://doi.org/10.1007/s12600-019-00734-1

Terekhova VV, Skrylnik YY (2012) Biological peculiarities of the alien for Europe *Anisandrus maiche* Stark (Coleoptera: Curculionidae: Scolytinae) bark beetle in Ukraine. Russian Journal of Biological Invasions 3(2): 139–144. https://doi.org/10.1134/S2075111712020105

Thurston GS, Slater A, Nei I, Roberts J, McLachlan Hamilton K, Sweeney JD, Kimoto T (2022) New Canadian and provincial records of Coleoptera resulting from annual Canadian Food Inspection Agency surveillance for detection of non-native, potentially invasive forest insects. Insects 13: e708. https://doi.org/10.3390/insects13080708

Tobin KN, Ginzel MD (2023) The ambrosia beetle *Anisandrus maiche* (Stark) is repelled by conophthorin and verbenone and attracted to ethanol in a dose-dependent manner. Agricultural and Forest Entomology 25: 103–110. https://doi.org/10.1111/afe.12534

Turner RM, Brockerhoff EG, Bertelsmeier C, Blake RE, Caton B, James A, MacLeod A, Nahrung HF, Pawson SM, Plank MJ, Pureswaran DS (2021) Worldwide border interceptions provide a window into human-mediated global insect movement. Ecological Applications 31: e02412. https://doi.org/10.1002/eap.2412

Vitale S, Santori A, Wajnberg E, Castagnone-Sereno P, Luongo L, Belisario A (2011) Morphological and molecular analysis of *Fusarium lateritium*, the cause of gray necrosis of hazelnut fruit in Italy. Phytopathology 101: 679–686. https://doi.org/10.1094/PHYTO-04-10-0120

White TJ, Bruns T, Lee S (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR Protocols: A Guide to Methods and Applications. Academic Press, Orlando, 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Supplementary material 1

Specimens of *Anisandrus maiche* from Switzerland, collected by José P. Ribeiro-Correia

Authors: José P. Ribeiro-Correia, Simone Prospero, Ludwig Beenken, Peter H. W. Biedermann, Simon Blaser, Manuela Branco, Yannick Chittaro, David Frey, Doris Hölling, Sezer Olivia Kaya, Miloš Knížek, Jana Mittelstrass, Beat Ruffner, Andreas Sanchez, Eckehard G. Brockerhoff

Data type: docx

Explanation note: Specimens of *Anisandrus maiche* from Switzerland, collected by José P. Ribeiro-Correia, caught in bottle traps with ethanol as lure, with information on collection locations and where specimens are held. See the manuscript for explanations of acronyms.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/alpento.8.117537.suppl1

Supplementary material 2

Specimens of *Anisandrus maiche* from Switzerland, collected by David Frey

Authors: José P. Ribeiro-Correia, Simone Prospero, Ludwig Beenken, Peter H. W. Biedermann, Simon Blaser, Manuela Branco, Yannick Chittaro, David Frey, Doris Hölling, Sezer Olivia Kaya, Miloš Knížek, Jana Mittelstrass, Beat Ruffner, Andreas Sanchez, Eckehard G. Brockerhoff

Data type: docx

Explanation note: Specimens of *Anisandrus maiche* from Switzerland, collected by David Frey, caught with unbaited Polytrap interception traps, with information on collection locations and where specimens are held. See the manuscript for explanations of acronyms.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/alpento.8.117537.suppl2

Supplementary material 3

Photos of traps used in this study

Authors: José P. Ribeiro-Correia, Simone Prospero, Ludwig Beenken, Peter H. W. Biedermann, Simon Blaser, Manuela Branco, Yannick Chittaro, David Frey, Doris Hölling, Sezer Olivia Kaya, Miloš Knížek, Jana Mittelstrass, Beat Ruffner, Andreas Sanchez, Eckehard G. Brockerhoff

Data type: docx

Explanation note: Traps used in Switzerland with which *Anisandrus maiche* was caught.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/alpento.8.117537.suppl3